
Aerial
United Kingdom • England • Wilts • Calstone Wellington
Buttercups Kindergarten
11. Look at soil compaction and nutrients to enhance sward growth rates
12. Consider gapping-up hedgerows to increase connectivity for wildlife
10. Redirect rainwater from cattle shed into drains or rainwater harvesting
13. Exclude wet area / flow pathway to capture flow and reduce sediment loading from livestock.
1. Consider moving current scraped yard material (slurry and manure) away from the river and move here.
9. Explore option for improved fish passage / connectivity (longer term)
14. Fencing to exclude livestock and create wider bufferstip (Explore CSF)
8. Riparian tree planting
2. Tree planting will help balance air quality emissions
7. Remove livestock drink
3. Move livestock Drink & feeder away from river and further into field
4. River restoration Site 2 (approx. 200m reach: Installation of LWD & strategic canopy thinning upstream
5. Explore options to create wider bufferstrip (tree planting & riparian fencing). Close off / remove existing livestock drink
6. Explore options to reduce impact of Spring runoff transporting farm runoff into the river (issue will be space due to pipework)
15. River restoration Site 1 (approx. 200m reach: Installation of LWD & strategic canopy thinning
16. Creation of bufferstrip incl. fencing & tree planting (different landowner). Exclude sheep from watercourse & reduce bank erosion
Feedback
100 feet
25 m
Image courtesy of Ordnance Survey, © 2022 TomTom